# <u>Chemistry 11 – Course Review</u>

### Introduction to Chemistry

1.  $0.0006 \text{ mm} = ? \mu \text{m}$ 

2. 0.054 mL = ? nL

Answer \_\_\_\_\_

Answer

3.  $3.5 \,\mu g/L = ? \,mg/mL$ 

Answer

4. The density of iron is 7860 g/L. Calculate the mass of a 3.2 mL sample of iron.

Answer

\_\_\_\_\_

5. Manganese has a density of 7.20 g/mL. Calculate the volume occupied by a 4.0 kg piece of manganese.


Answer \_\_\_\_\_


Name:	Block:	Date:
Chemistry 11	Organic Chemistry Review	Assignment


1. Identify the following as either an alkane, alkene or alkyne.

a. C<sub>4</sub>H<sub>8</sub> b. C<sub>4</sub>H<sub>10</sub> c. C<sub>14</sub>H<sub>26</sub> d. C<sub>40</sub>H<sub>78</sub> e. C<sub>25</sub>H<sub>52</sub>

2.Identify which class of organic compounds each of the following belongs to: halocarbon, alcohol, ether, aldehyde, ketone, carboxylic acid, ester, amine, amide, alkane, alkene, or alkyne. Name each compound.







# 3. Draw the following organic compounds. Identify what class or organic compounds it belongs to.

- a. decanoic acid
- b. methyl octanoate
- c. 5-ethyl-2-heptyne
- d. 3,3,4,5-tetraiodo-1-heptene
- e. 3-methyl-1-butyne
- f. 4,4-diethyl-1-hexyne
- g. 2,2-dimethylpentanoic acid
- h. methylpropylamine
- i. 3-methyl-2-pentene
- j. pentyl butanoate
- k.2-pentanone
- I. 3-ethyl-4,5-dipropyloctane

- m. 4-ethyl-5-methyl-2-hexanol
- n. 2,3-dimethylbutanal
- o. 1-butyl-2,3-dimethylcyclopentane
- p. 1,3-dichloro-2-fluorocyclopentane
- q. 3-methylbutanamide
- r. 4-isopropyldecane
- s. 3-ethoxypentane
- t. 6-isopropyl-2,8-dimethylnonene
- u. 2-ethyl-3-methylpentanal
- v. 1-methoxybutane
- w. 3-methyl-2-butanone
- x. 3,4-dimethylcylcopentanol

6. A 0.0460 L piece of copper has a mass of 410.32 g. Calculate the density of copper in g/mL.

		21 14		Answer	
7.		ve the number of significant digits in each asurements.	ch o	f the following. Assume they are all	Sa <sub>pela</sub>
	a)	0.0023	d)	3.2 x 10 <sup>-4</sup>	
ос. Ю	b)	3953 000	e)	50020.000	
	c)	1.0200 x 10 <sup>5</sup>	f)	3450	(e) - A)
8.		rform the following calculations and ro gnificant digits as justified by the data. A		the answers off to the correct number of ime all numbers are measurements.	В 6 <sup>гл</sup>
	a)	2.1500 x 0.31	f)	$8.90 \times 10^3 \div 4.400 \times 10^{-6}$	
÷	b)	0.05 + 394.7322	g)	$83.00 \div 1.2300 \times 10^2$	,
	c)	$4.905 \ge 10^6 \div 4 \ge 10^{-2} \dots$	h)	98.0076 - 2.195	<u> </u>
	d)	(3.33 x 9.52) + 13.983	i)	0.00000200 x 245.912	α
	e)	3.813 + 98.98 + 2.669	j)	5.802 ÷ 6.21 + 2.41 ÷ 9.2565	
9.	Ro	ound the following numbers to 2 signific	cant	digits. (4 marks)	
	a)	2 000 000 000	c)	3.88945 x 10 <sup>28</sup>	
	b)	106 000	d)	0.000 000 7895	
		v		14) 14)	

### Properties of Matter

2.

1. Define: Observation, Interpretation, Qualitative, Quantitative, Data, Experiment, Hypothesis, Theory, Laws, Matter, Chemistry, Physical and Chemical Properties, Malleability, Ductility, Lustre, Viscosity and Diffusion. Review the Phases of Matter.

Draw the diagram from your notes outlining the Classification of Matter. Make sure you can define each classification.

- 3. Concerning separation techniques...
  - a) Explain how distillation can be used to separate the substances in a solution.


- b)
- What types of mixtures does paper chromatography work best for?
- c) Explain how a centrifuge separates the components of a suspension.
- 4. Define a physical change –

Give some examples of physical changes.

5. Define a chemical change –

Give some examples of chemical changes.

Given the following graph of Temperature vs. Time for warming substance "X" which starts out as a solid, answer the questions below:



6.

Na	mes	and Formulas for Compounds			х — Ц (20)	
1.	Wri	te the correct formula for the following compounds:	e 8	$x^*$	3	
10	a)	ammonium chlorate	а а		* 	
	b)	copper (II) sulphite				4
	c)	zinc carbonate tetrahydrate	(		5	
	d)	nitric acid				
	e)	phosphorus pentaiodide		X		
2	f)	iron (III) thiocyanate				34 
	g)	sulphuric acid	5 	13		
	h)	dinitrogen tetrafluoride	53 		2	
2.	Wri	te the correct names for the following compounds:	e.	( P		
	a)	Mn(SO <sub>4</sub> ) <sub>2</sub>				ă.
	b)	PbCrO <sub>4</sub> <sup>•</sup> 6H <sub>2</sub> O				
	c)	As <sub>2</sub> O <sub>3</sub>				
2	d)	СН3СООН				acid
	e)	Ni <sub>2</sub> (C <sub>2</sub> O <sub>4</sub> ) <sub>3</sub>	e 8			
	f)	NF3				
	g)	(NH <sub>4</sub> ) <sub>2</sub> HPO <sub>4</sub>			<u>.</u>	
	g) h)	$(NH_4)_2HPO_4$ Ba $(OH)_2$ ·10H <sub>2</sub> O		ş	<u>, -</u>	

### The Mole Concept

1. Make the following conversions, clearly showing your steps. Include proper units in all of your work and in your answer.

Answer

Answer \_\_\_\_

Answer

a) 133.44 grams of  $PCl_5 = ?$  moles

b) 0.00256 moles of  $Li_2Cr_2O_7 = ?$  grams

c)  $170.24 \text{ L of NO}_2 \text{ at STP} = ? \text{ moles}$ 

d) 570.625 g of PCl<sub>3</sub> gas = ? L (STP)

1030.4 mL of  $C_2H_6$  gas at STP = ? g

e)

Answer\_\_\_\_\_

\_\_\_\_\_

Answer \_\_\_\_\_

f) 5.00 kg of nitrogen gas = ? L (STP)

g) 0.5696 kg of  $CH_{4(g)} = ? mL (STP)$ 

 The density of liquid ethanol (C<sub>2</sub>H<sub>5</sub>OH) is 0.790 g/mL. Calculate the number of molecules in a 35.0 mL sample of liquid ethanol. (NOTE: You CAN'T use 22.4 L/mol since this is NOT a gas at STP!)

Answer

\_\_\_\_\_

Answer

Answer

3. A 100.0 mL sample of liquid mercury contains 6.78 moles. Calculate the density of liquid mercury from this data.

4. Calculate the density of  $PCl_{3(g)}$  at STP.

Answer

Answer

- a) The density of a gas at STP is 4.955 g/L. Calculate the molar mass of this gas.
  - b) The gas is an oxide of selenium. Determine the molecular formula.

5.

6. Find the percent composition (% by mass of each element) in the following compound: Sr<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>. Show your work.

Answer

Answer

%Sr.

A compound was analyzed and the following results were obtained:
Molar mass: 270.4 g/mol
Mass of sample: 162.24 g
Mass of potassium: 46.92 g
Mass of sulphur: 38.52 g
Mass of oxygen: the remainder of the sample is oxygen

a) Determine the mass of oxygen in the sample.

Answer

%P.

%O

b) Determine the empirical formula for this compound.

Answer: Empirical Formula:

c) Determine the molecular formula for this compound.

Answer: Molecular Formula:

8. 123.11 g of zinc nitrate, Zn(NO<sub>3</sub>)<sub>2</sub> are dissolved in enough water to form 650.0 mL of solution. Calculate the [Zn(NO<sub>3</sub>)<sub>2</sub>]) Include proper units in your work and in your answers.

Answer

9. Calculate the mass of potassium sulphite (K<sub>2</sub>SO<sub>3</sub>) needed to make 800.0 mL of a 0.200 M solution of K<sub>2</sub>SO<sub>3</sub>. Include proper units in your work and in your answers.

Answer

10. What volume of 2.50 M Li<sub>2</sub>CO<sub>3</sub> would need to be evaporated in order to obtain 47.232 g of solid Li<sub>2</sub>CO<sub>3</sub>? Include proper units in your work and in your answers.

Answer

11. 150.0 mL of water are added to 400.0 mL of 0.45 M HNO<sub>3</sub>. Calculate the final [HNO<sub>3</sub>]. Include proper units in your work and in your answers.

Answer

12. What volume of water needs to be added to 150.0 mL of 4.00 M H<sub>2</sub>SO<sub>4</sub> in order to bring the concentration down to 2.50 M? Include proper units in your work and in your answers.

Answer

13. Give directions on how to make 5.00 L of 0.020 M Ca(ClO)<sub>2</sub> using solid Ca(ClO)<sub>2</sub> and water. Include proper units in your work and in your answers.

Directions:

A) Write the dissociation reaction and then Calculate the concentration of each ion in the following solutions.

- 1) 0.50 M HCI
- 2) 3.00 M H<sub>2</sub>SO<sub>4</sub>
- 3) 6.42 g Na<sub>3</sub>PO<sub>4</sub> dissolved in water to make 25.5 mL of solution.
- 4) 1.00 g CuSO<sub>4</sub>·5H<sub>2</sub>O dissolved in water to make 50.0 mL of solution.

#### B) Calculate

- a) how many moles of solute
- b) how many moles of each ion are in the following solutions.
- 5) 25.0 mL of 2.50 M NaOH
- 6) 10.0 mL of 0.0500 M CoCl<sub>2</sub>
- 7) 117 mL of 14.6 M H3PO4
- 8) 11.3 mL of 0.512 M Ca(OH)<sub>2</sub>

#### **C) Dilution Problems**

Calculate the concentration of a) each solute and b) each ion when the following are mixed.

- 9) 25.0 mL of 18.0 M H<sub>2</sub>SO<sub>4</sub> and 130.0 mL water
- 10) 110.0 mL of 3.00 M HCl and 25.0 mL water
- 11) 50.0 mL of 0.750 M KOH and 25.0 mL of 0.500 M KOH
- 12) 75.0 mL of 1.00 M LiNO3 and 115 mL of 2.50 M FeCl3

C) Some chemicals come from suppliers as highly concentrated solutions. In order to use them properly (and safely) they need to be diluted. Describe how you would make the following solutions given the strength of the initial solution. (State the volume of the stock solution needed.)

13) 250.0 mL of 3.00 M HCl using 12.0 M HCl
14) 50.0 mL of 1.00 M H<sub>2</sub>SO<sub>4</sub> using 18.0 M H<sub>2</sub>SO<sub>4</sub>
15) 100.0 mL of 0.500 M CH<sub>3</sub>COOH using 15.0 M CH<sub>3</sub>COOH

16) 500.0 mL of 1.50 M NH<sub>3</sub> using 15.0 M NH<sub>3</sub>

#### **Chemical Reactions**

1. Balance the following equations

 $NH_3 + O_2 \rightarrow NO + H_2O$ 

 $(NH_4)_2C_2O_4 + AlCl_3 \rightarrow Al_2(C_2O_4)_3 + NH_4Cl$ 

 $C_{14}H_{30} + O_2 \rightarrow CO_2 + H_2O$ 

Fe + HNO<sub>3</sub>  $\rightarrow$  Fe(NO<sub>3</sub>)<sub>3</sub> + H<sub>2</sub>

 $P_4 + Cl_2 \rightarrow PCl_3$ 

 $Na_2Cr_2O_7 + HCl \rightarrow NaCl + CrCl_3 + H_2O + Cl_2$ 

 $H_3PO_4 + Ca(OH)_2 \rightarrow Ca_3(PO_4)_2 + H_2O$ 

 $Ba(ClO_4)_2 \rightarrow Ba + Cl_2 + O_2$ 

 $C_7H_{15}OH + O_2 \rightarrow CO_2 + H_2O$ 

 $MgSO_4 5H_2O \rightarrow MgSO_4 + H_2O$ 

- 2. Write a balanced chemical equation for each of the following, and classify each as synthesis, decomposition, single replacement, double replacement, neutralization or combustion.
  - a) potassium sulphate is mixed with cobalt (III) nitrate
  - b) liquid propanol ( $C_3H_7OH$ ) is burned in air

c) ammonium nitrate is decomposed into it's elements

- d) a piece of zinc is placed in a test-tube containing a solution of silver nitrate
- e) bromine reacts with sodium iodide
- f) bromine reacts with aluminum
- g) rubidium reacts with chlorine gas
- h) hydrochloric acid reacts with strontium hydroxide
- 3. State whether each of the following are *exothermic* or *endothermic*.

$HCl + 432 \text{ kJ} \rightarrow H + Cl$	Answer
$C_{12}H_{22}O_{11} + 12 O_2 \rightarrow 12CO_2 + 11H_2O$	$\Delta H = -5638 \text{ kJ}$ Answer
$H_2O_{(s)} \rightarrow H_2O_{(l)}$	Answer
Ellergy (kJ)	ž.
Reaction Proceeding $\rightarrow$	
	Answer
$CD \rightarrow C + D$ $\Delta H = 65.7 \text{ kJ}$	Answer
$E + F + 437 \text{ kJ} \rightarrow G + H$	Answer

Page 13

### Given the equation: $C_{12}H_{22}O_{11} + 12O_2 \rightarrow 12CO_2 + 11H_2O + 5638 \text{ kJ}$

4.

c.

a. How much heat is released during the formation of  $880.0 \text{ g of } \text{CO}_2$ ?

Answer

b. How much heat is released during the formation of 5.6 moles of  $H_2O$ ?

Answer

If 179.2 L of O<sub>2</sub> (STP) are consumed, how much heat is released?

Answer

### Stoichiometry

1. Given the following balanced equation, answer the questions following it:

 $2NF_{3(g)} + 3H_{2(g)} \rightarrow N_{2(g)} + 6HF_{(g)}$ 

a) If 5.5 moles of  $H_2$  are reacted, how many moles of NF<sub>3</sub> will be consumed?

Answer

b) In order to produce 0.47 moles of HF, how many moles of NF<sub>3</sub> would be consumed?

Answer \_\_\_\_\_

c) If you needed to produce 180.6 g of  $N_2$ , how many moles of  $H_2$  would you need to start with?

Answer \_\_\_\_\_

d) If you completely react 17.04 g of NF<sub>3</sub>, what mass of HF will be produced?

Answer \_\_\_\_

2. Given the following balanced equation, answer the questions following it:

HBrO<sub>3</sub> + 5 HBr  $\rightarrow$  3 H<sub>2</sub>O<sub>(1)</sub> + 3 Br<sub>2(g)</sub>

- a) If 3.56 moles of HBr are reacted, how many Litres of Br<sub>2</sub> will be formed at STP?
- b) In order to produce  $3.311 \times 10^{24}$  molecules of Br<sub>2</sub>, what mass of HBr is needed?

#### Answer

Answer

3. Given the following balanced chemical equation, answer the question below it.

 $MgCO_{3(s)} + 2HCl_{(aq)} \rightarrow CO_{2(g)} + H_2O_{(l)} + MgCl_{2(aq)}$ 

a) What mass of MgCO<sub>3</sub> will react completely with 15.0 mL of 1.5 M HCl?

Answer

b) Calculate the volume of 2.0 M HCl which would be needed to react completely with 37.935 grams of magnesium carbonate.

Answer \_\_\_\_\_

4. Given the following balanced equation, answer the questions below it.

 $Ba(OH)_{2(aq)} + 2 HNO_{3(aq)} \rightarrow 2 H_2O_{(1)} + Ba(NO_3)_2$ 

a) In a titration, 18.20 mL of 0.300 M Ba(OH)<sub>2</sub> is required to react completely with a 25.0 mL sample of a solution of HNO<sub>3</sub>. Find the [HNO<sub>3</sub>].

#### Answer \_

b) In a titration, 11.06 mL of  $0.200 \text{ M HNO}_3$  is required to react completely with a sample of  $0.250 \text{ M Ba}(\text{OH})_2$ . Find the volume of the Ba(OH)<sub>2</sub> sample.

Answer

5. Given the following balanced equation, answer the questions below it.

 $3 \operatorname{Cu}_{(s)} + 8 \operatorname{HNO}_{3(l)} \rightarrow 3 \operatorname{Cu}(\operatorname{NO}_3)_{2(aq)} + 2 \operatorname{NO}_{(g)} + 4 \operatorname{H}_2 O_{(l)}$ 

a)

If 317.5 grams of Cu are placed into 756.0 grams of  $HNO_3$ , determine which reactant is in excess.

Answer \_\_\_\_\_

b) If the reaction in (a) is carried out, what mass of NO will be formed?

Answer

6. Given the balanced equation:  $2BN + 3F_2 \rightarrow 2BF_3 + N_2$ , When 161.2 grams of BN are added to an excess of F<sub>2</sub>, a reaction occurs in which 326.118 grams of BF<sub>3</sub> are formed.

a) Calculate the *theoretical* yield of BF<sub>3</sub> in grams.

b) Calculate the *percentage* yield of BF<sub>3</sub>.

Answer

Answer

Answer

Answer

7. When reacting  $NH_3$  with  $O_2$  according to the reaction:

 $4 \text{ NH}_3 + 5 \text{ O}_2 \rightarrow 4 \text{ NO} + 6 \text{ H}_2\text{O}$ 

Using 163.2 grams of  $NH_3$  with an excess of  $O_2$  produces a 67% yield of NO.

a) Calculate the *theoretical yield* of NO in grams.

b) Calculate the *actual yield* of NO in grams.

#### Atoms, Periodic Table and Bonding

2. Consider the following ideas:

3.

4.

- > Compounds are made up of molecules which are combinations of atoms
- > All atoms of an element are the same
- Atoms of different elements are different
- > Atoms are indivisible particles

Who came up with these ideas? \_\_\_\_\_\_ He called the ideas, the

Theory.

\_\_\_\_\_ measured the charge/mass ratio of an electron and came up with the so-called "plum pudding" model of the atom.

\_\_\_\_\_\_ devised the Scattering Experiment, which showed that all atoms had a small dense \_\_\_\_\_\_.

5. Bohr came up with an atomic model to explain the spectrum of

He said that the atom has certain \_\_\_\_\_\_ levels which are allowed. These levels corresponded to \_\_\_\_\_\_ in which electrons move. If an electron absorbs a certain photon of energy, it will jump to a \_\_\_\_\_\_ level. It will release this energy (in the form of \_\_\_\_\_\_) when it jumps back to a \_\_\_\_\_\_ level.

What were two limitations of Bohr's atomic model?

Isotope	Protons	Neutrons	Electrons
$^{194}$ Ir <sup>3+</sup>	à.	1. E	
$^{202}\text{Hg}^{2+}$			
$^{125}$ Te <sup>2-</sup>			
<sup>263</sup> Sg	1	4	
$^{2}\text{H}^{+}$			

#### 6. Give the number of protons, neutrons and electrons in the following:

7. Give the nuclear notation of the following:

Isotope	Protons	Neutrons	Electrons
-to stating the set	105	157	103
12	51	72	48
	33	42	36
50 a 1	54	79	54
<u> (18</u> -11-17-17-17-17-17-17-17-17-17-17-17-17-	94	150	91

8. Element "X" is composed of the following naturally occurring isotopes:

Isotope	% Abundance
<sup>79</sup> X	50.69
X <sup>18</sup>	49.31

Calculate the average atomic mass of element "X" to 3 decimal places.

Element "X" is actually the real element

9. Regions in space occupied by electrons are called

10. The principal quantum number is given the letter \_\_\_\_\_ and refers to the \_\_\_\_\_\_ level.

11.

Write the ground state electron configurations (eg.  $1s^2 2s^2 2p^6$ ) for the following atoms or ions. You may use the core notation.

- a) P
- b) Mo
- c) Se
- d) Rb
- e) Cl
- f) Al<sup>3+</sup>
- g) K<sup>+</sup>
- h) S<sup>2-</sup>

13.

14.

12. In order to become stable,

an atom of Sr will	electrons and become the ion
an atom of As will	electrons and become the ion
an atom of Al will	electrons and become the ion
an atom of Se will	electrons and become the ion
an atom of N will	electrons and become the ion
an atom of I will	_ electrons and become the ion
an atom of Cs will	electrons and become the ion
an atom of Te will	electrons and become the ion
Circle the metalloid: Be Rb O	s Ge Pb Al
Circle the most reactive element in	n the following: Na Mg Si Al Ar

15. Circle the most reactive element in the following: Na K Rb Cs Li

16. Circle the most reactive element in the following: Cl Br I At Ne

- 17. Circle the element with the largest atomic radius of these: Na Mg Si Al Ar
- 18. Circle the element with the largest atomic radius of these: N P As Sb Bi

19.	Circle the element with the largest ionization energy of these: K Ca Ga As Kr
20.	Circle the element with the largest ionization energy of these: C Si Ge Sn Pb
21.	What is meant by ionization energy?
22.	Circle the element with the largest density of these: C Si Ge Sn Pb
23.	Circle the element with the largest density of these: Na K Rb Cs Li
24.	Circle the element with the highest electronegativity of these: Mg Sr Ba Ra
25.	Circle the element with the highest electronegativity of these: Mg Si S Cl
26.	Circle the element with the highest electronegativity of these: F Cl Br I
27.	What is meant by electronegativity?
28.	Circle the most metallic element of these: Be Mg Ca Sr Ba
29.	Circle the most metallic element of these: B Al Ga In Tl
30.	Circle the most metallic element of these: Ga Ge Se Br Kr
31.	In an ionic bond, electrons are <ul> <li>a. shared equally by two atoms</li> <li>b. shared unequally by two atoms</li> <li>c. transferred from a metal to a non-metal</li> <li>d. transferred from a non-metal to a metal</li> <li>e. closer to one end of a molecule, forming a temporary dipole Answer</li></ul>
32.	In a covalent bond, electrons are f. shared equally by two atoms g. shared unequally by two atoms h. transferred from a metal to a non-metal i. transferred from a non-metal to a metal j. closer to one end of a molecule, forming a temporary dipole Answer
33.	<ul> <li>In a polar covalent bond, electrons are</li> <li>k. shared equally by two atoms</li> <li>l. shared unequally by two atoms</li> <li>m. transferred from a metal to a non-metal</li> <li>n. transferred from a non-metal to a metal</li> <li>o. closer to one end of a molecule, forming a temporary dipole</li> </ul>

# Page 22

 34. In London forces, electrons are

- p. shared equally by two atoms
- q. shared unequally by two atoms
- r. transferred from a metal to a non-metal
- s. transferred from a non-metal to a metal
- t. closer to one end of a molecule, forming a temporary dipole

Answer

35. What evidence do we have that ionic bonds are very strong?

36. Write electron-dot diagrams for: MgCl<sub>2</sub> (ionic) PBr<sub>3</sub>(covalent)

 $SeF_2(covalent)$ 

CH<sub>3</sub>CH<sub>2</sub>I(covalent)

## Yet More Lewis Structures

For those of you that enjoy such things, some more Lewis structures to draw:

6.)  $P_{4}^{-3}$ BSF

3)  $C_2H_5OH$  (ethanol)

1)

2)

HBr

5) SF<sub>6</sub>

 $N_2F_4$ 

4)

For chemistry help, visit www.chemfiesta.com