

Chemistry 11 – Course Review

KEY

Unit 2—Introduction to Chemistry

1. $0.0006 \text{ mm} = ? \mu\text{m}$

$$6 \times 10^{-4} \text{ mm} \times \frac{10^{-3} \text{ m}}{1 \text{ mm}} \times \frac{1 \mu\text{m}}{10^{-6} \text{ m}}$$

Answer 0.6 μm

2. $0.054 \text{ mL} = ? \text{ nL}$

$$5.4 \times 10^{-2} \text{ mL} \times \frac{10^{-3} \text{ L}}{1 \text{ mL}} \times \frac{1 \text{ nL}}{10^{-9} \text{ L}}$$

Answer $5.4 \times 10^4 \text{ nL}$

3. $3.5 \mu\text{g/L} = ? \text{ mg/mL}$

$$\frac{3.5 \mu\text{g}}{\text{L}} \times \frac{10^{-6} \text{ g}}{1 \mu\text{g}} \times \frac{1 \text{ mg}}{10^{-3} \text{ g}} \times \frac{10^{-3} \text{ L}}{1 \text{ mL}}$$

Answer $3.5 \times 10^{-6} \text{ mg/mL}$

4. The density of iron is 7860 g/L . Calculate the mass of a 3.2 mL sample of iron.

$$0.0032 \text{ L} \times \frac{7860 \text{ g}}{\text{L}}$$

Answer 25.2g

5. Manganese has a density of 7.20 g/mL . Calculate the volume occupied by a 4.0 kg piece of manganese.

$$V = \frac{m}{D} = \frac{4000 \text{ g}}{7.20 \text{ g/mL}} = 555.56$$

Answer 555 mL

6. A 0.0460 L piece of copper has a mass of 410.32 g . Calculate the density of copper in g/mL .

$$D = \frac{m}{V} = \frac{410.32 \text{ g}}{46 \text{ mL}} = 8.92 \text{ g/mL}$$

Answer _____

7. Give the number of significant digits in each of the following. Assume they are all measurements.

- a) 0.0023 2 d) 3.2×10^{-4} 2
 b) 3953 000 4 e) 50020.000 8
 c) 1.0200×10^5 5 f) 3450 3

8. Perform the following calculations and round the answers off to the correct number of significant digits as justified by the data. Assume all numbers are measurements.


- a) 2.1500×0.31 0.67 f) $8.90 \times 10^3 \div 4.400 \times 10^6$ 2.02×10^{-7}
 b) $0.05 + 394.7322$ 394.78 g) $83.00 \div 1.2300 \times 10^2$ 0.6748
 c) $4.905 \times 10^6 \div 4 \times 10^{-2}$ 1×10^8 h) $98.0076 - 2.195$ 95.813
 d) $(3.33 \times 9.52) + 13.983$ 45.7 i) $0.00000200 \times 245.912$ ~~###~~ 4.92×10^{-7}
 e) $3.813 + 98.98 + 2.669$ 105.46 j) $5.802 \div 6.21 + 2.41 \div 9.2565$ 1.195

9. Round the following numbers to 2 significant digits. (4 marks)

- a) 2 000 000 000 2.0×10^9 c) 3.88945×10^{28} 3.9×10^{28}
 b) 106 000 1.1×10^5 d) 0.000 000 7895 7.9×10^{-7}

Unit 3—Properties of Matter

1 Draw the diagram from your notes outlining the Classification of Matter. Make sure you can define each classification.



ANSWER KEY



















1. Identify the following as either an alkane, alkene or alkyne.







- a. alkene b. alkane c. alkyne d. alkyne e. alkane

2. Identify which class of organic compounds each of the following belongs to: halocarbon, alcohol, ether, aldehyde, ketone, carboxylic acid, ester, amine, amide, alkane, alkene, or alkyne. Name each compound.

- a. **Ketone; 4,4-dimethyl-2-pentanone**
- b. **Halocarbon; 3,3-dibromo-2,2-dichlorohexane**
- c. **Alkyne; 4-ethyl-7-methyl-2-octyne**
- d. **Alkane; 3,4-dimethylhexane**
- e. **Carboxylic acid; methylpropanoic acid**
- f. **Ester; methylhexanoate**
- g. **Alkane; 7-ethyl-3-methyldecane**
- h. **Alkene; 4,4,-dimethyl-1-pentene**
- i. **Ketone; propanone**
- j. **Alcohol; 2-pentanol**
- k. **Ester; ethylbutanoate**
- l. **Carboxylic acid; propanoic acid**
- m. **Aldehyde; propanal**
- n. **Aldehyde; 2-methylbutanal**
- o. **Cyclic hydrocarbon; 1,3,-diethyl-6,8-dimethylcyclononane**
- p. **Cyclic halocarbon; 1,4-dichloro-2,3-diethylcyclopentane**
- q. **Amine; trimethylamine**
- r. **Amide; propanamide**
- s. **Ether; 1-propoxypropane**

3. ANSWERS ON THE FOLLOWING PAGE





2. Define a physical change - Change where chemical make up does not change
Give some examples of physical changes.


Rip paper

3. Define a chemical change -
new substance is formed

Give some examples of chemical changes.

burning ; cooking ; neutralization

4. Given the following graph of Temperature vs. Time for warming substance "X" which starts out as a solid, answer the questions below:



- a) During time 0.0 - 5.0 minutes, the added heat energy is being used to incr. temp of solid
- b) During time 5.0 - 15.0 minutes, the added heat energy is being used to break bonds holding together solid
- c) During time 15.0 - 20.0 minutes, the added heat energy is being used to increase T of liquid
- d) During time 20.0 - 28.0 minutes, the added heat energy is being used to break bonds of liquid
- e) The melting point of substance "X" is 43 °C
- f) The boiling point of substance "X" is 77 °C
- g) If a greater amount of substance "X" was used, the melting point would be
1. a lower temperature
2. a higher temperature
3. the same temperature Answer 3
- h) What phase is substance "X" at 90°C? Gas

Unit 4— Names and Formulas for Compounds

1. Write the correct formula for the following compounds:

- a) ammonium chlorate NH_4ClO_3
- b) copper (II) sulphite..... CuSO_3
- c) zinc carbonate tetrahydrate $\text{ZnCO}_3 \cdot 4\text{H}_2\text{O}$
- d) nitric acid HNO_3
- e) phosphorus pentaiodide PI_5
- f) iron (III) thiocyanate..... $\text{Fe}(\text{SCN})_3$
- g) sulphuric acid..... H_2SO_4
- h) dinitrogen tetrafluoride N_2F_4

2. Write the correct names for the following compounds:

- a) $\text{Mn}(\text{SO}_4)_2$ Manganese (IV) sulphate
- b) $\text{PbCrO}_4 \cdot 6\text{H}_2\text{O}$ lead (II) chromate hexahydrate
- c) As_2O_3 diarsenic trioxide
- d) CH_3COOH Acetic Acid acid
- e) $\text{Ni}_2(\text{C}_2\text{O}_4)_3$ nickel (III) oxalate
- f) NF_3 Nitrogen trifluoride
- g) $(\text{NH}_4)_2\text{HPO}_4$ Ammonium monohydrogen phosphate
- h) $\text{Ba}(\text{OH})_2 \cdot 10\text{H}_2\text{O}$ Barium hydroxide decahydrate

Unit 5— The Mole Concept

1. Make the following conversions, clearly showing your steps. Include proper units in all of your work and in your answer.

- a) 133.44 grams of PCl_5 = ? moles

$$133.44 \text{ g} \times \frac{1 \text{ mol}}{208.5 \text{ g}}$$

Answer 0.64 mol

d) 570.625 g of PCl_3 gas = ? L (STP)

$$570.625 \text{ g} \times \frac{1 \text{ mol}}{137.5 \text{ g}} \times \frac{22.4 \text{ L}}{1 \text{ mol}} =$$

Answer 92.96 L

e) 1030.4 mL of C_2H_6 gas at STP = ? g

$$1.0304 \text{ L} \times \frac{1 \text{ mol}}{22.4 \text{ L}} \times \frac{30.0 \text{ g}}{1 \text{ mol}}$$

Answer 1.38

f) 5.00 kg of nitrogen gas = ? L (STP)

$$5000 \text{ g} \times \frac{1 \text{ mol}}{28.0 \text{ g}} \times \frac{22.4 \text{ L}}{1 \text{ mol}}$$

Answer 4000. L

g) 0.5696 kg of $\text{CH}_4(\text{g})$ = ? mL

$$0.5696 \text{ kg} = 569.6 \text{ g} \times \frac{1 \text{ mol}}{16.0 \text{ g}} \times \frac{22.4 \text{ L}}{1 \text{ mol}} \times \frac{1 \text{ mL}}{10^{-3} \text{ L}}$$

Answer $7.97 \times 10^5 \text{ mL}$

2. The density of liquid ethanol ($\text{C}_2\text{H}_5\text{OH}$) is 0.790 g/mL. Calculate the number of molecules in a 35.0 mL sample of liquid ethanol. (NOTE: You CAN'T use 22.4 L/mol since this is NOT a gas at STP!)

$$\frac{0.790 \text{ g}}{\text{mL}} \times 35 \text{ mL} = 27.65 \text{ g} \times \frac{1 \text{ mol}}{46 \text{ g}} \times \frac{6.022 \times 10^{23} \text{ molec}}{1 \text{ mol}}$$

Answer

$= 3.62 \times 10^{23} \text{ molec.}$

7. A compound was analyzed and the following results were obtained:

Molar mass: 270.4 g/mol

Mass of sample: 162.24 g

Mass of potassium: 46.92 g

Mass of sulphur: 38.52 g

Mass of oxygen: the remainder of the sample is oxygen

a) Determine the mass of oxygen in the sample.

$$162.24\text{g} - 46.92 - 38.52$$

Answer 76.8

b) Determine the empirical formula for this compound.

$$\text{K: } \frac{46.92}{162.24} \times 100\% \times \frac{1\text{mol}}{39.1\text{g}} = 1.20 \div 1.2 = 1$$

$$\text{S: } \frac{38.52}{162.24} \times 100\% \times \frac{1\text{mol}}{32.1\text{g}} = 1.2 \div 1.2 = 1$$

$$\text{O: } \frac{76.8}{162.24} \times 100\% \times \frac{1\text{mol}}{16\text{g}} = 4.8 \div 1.2 = 4$$

Answer: Empirical Formula: KSO₄

c) Determine the molecular formula for this compound.

$$\frac{\text{MF}}{\text{EF}} = \frac{270.4\text{g/mol}}{135.2\text{g/mol}} = 2 \times \text{KSO}_4$$

Answer: Molecular Formula: K₂S₂O₈

8. 123.11 g of zinc nitrate, Zn(NO₃)₂ are dissolved in enough water to form 650.0 mL of solution. Calculate the [Zn(NO₃)₂] Include proper units in your work and in your answers.

$$123.11\text{g} \times \frac{1\text{mol}}{189.4\text{g}} = 0.65\text{mol}$$

$$[\text{Zn}(\text{NO}_3)_2] = \frac{0.65\text{mol}}{0.65\text{L}}$$

Answer 1.000M

9. Calculate the mass of potassium sulphite (K_2SO_3) needed to make 800.0 mL of a 0.200 M solution of K_2SO_3 . Include proper units in your work and in your answers.

$$\frac{0.200 \text{ mol}}{\cancel{L}} \times 0.800 \cancel{L} = 0.160 \text{ mol} \times \frac{158.3 \text{ g}}{1 \text{ mol}}$$

Answer 25.328g

10. What volume of 2.50 M Li_2CO_3 would need to be evaporated in order to obtain 47.232 g of solid Li_2CO_3 ? Include proper units in your work and in your answers.

$$47.232 \text{ g} \times \frac{1 \text{ mol}}{73.8 \text{ g}} = 0.64 \text{ mol}$$



$$V = \frac{0.64 \text{ mol}}{2.50 \text{ M}} =$$

Answer 0.256 L

11. 150.0 mL of water are added to 400.0 mL of 0.45 M HNO_3 . Calculate the final $[HNO_3]$. Include proper units in your work and in your answers.

$$m_1 v_1 = m_2 v_2$$

$$m_2 = \frac{(0.45 \text{ M})(400 \text{ mL})}{(500 \text{ mL})} =$$

Answer 0.327 M

12. What volume of water needs to be added to 150.0 mL of 4.00 M H_2SO_4 in order to bring the concentration down to 2.50 M? Include proper units in your work and in your answers.





$$m_1 v_1 = m_2 v_2$$






$$\frac{(4.00 \text{ M})(150 \text{ mL})}{(2.50 \text{ M})} \rightarrow v_2 = 240 \text{ mL} - 150 \text{ mL}$$

Answer 90 mL

Unit 6—Chemical Reactions


1. Balance the following equations






2. Write a balanced chemical equation for each of the following, and classify each as synthesis, decomposition, single replacement, double replacement, neutralization or combustion.


- a) potassium sulphate is mixed with cobalt (III) nitrate



- b) liquid propanol (C_3H_7OH) is burned in air




- c) ammonium nitrate is decomposed into its elements




- d) a piece of zinc is placed in a test-tube containing a solution of silver nitrate




- e) bromine reacts with sodium iodide




- f) bromine reacts with aluminum




- g) rubidium reacts with chlorine gas




- h) hydrochloric acid reacts with strontium hydroxide




3. State whether each of the following are *exothermic* or *endothermic*.




Answer endo



$\Delta H = -5638 \text{ kJ}$ Answer exo




Answer endo




exo.


Answer



Answer

endo

Answer

endoa. How much heat is released during the formation of 880.0 g of CO_2 ?

$$880.0 \times \frac{1 \text{ mol } CO_2}{44 \text{ g } CO_2} \times \frac{5638 \text{ kJ}}{12 \text{ mol}}$$

Answer

9396.67 kJb. How much heat is released during the formation of 5.6 moles of H_2O ?

$$5.6 \text{ mol } H_2O \times \frac{5638 \text{ kJ}}{12 \text{ mol } H_2O}$$

Answer

2870.25 kJ
~~3758.67 kJ~~c. If 179.2 L of O_2 (STP) are consumed, how much heat is released?~~XXXXXXXXXX~~

$$179.2 \text{ L} \times \frac{1 \text{ mol } O_2}{22.4 \text{ L}} \times \frac{5638 \text{ kJ}}{12 \text{ mol } O_2}$$

Answer

3758.67 kJ5. Calculate the amount of heat (in Joules) required to warm 200.0 g of water from 8.0°C to 45.0°C . (Heat Capacity (C) for H_2O is $4180 \text{ J/kg} \cdot ^\circ\text{C}$)

$$E = mc\Delta T$$
$$= (0.200 \text{ kg})(4180)(37^\circ\text{C})$$

=

Answer

30932 J6. 13.376 kJ of heat are added to a 400.0 gram sample of water initially at 4.0°C . Calculate the final temperature of the water sample. Be careful with units! (Heat Capacity (C) for H_2O is $4180 \text{ J/kg} \cdot ^\circ\text{C}$)

$$E = mc\Delta T$$

$$13376 \text{ J} = (0.4 \text{ kg})(4180)\Delta T$$

$$\Delta T = 8^\circ\text{C}$$

Answer

 12°C

$$\text{Final} = 4^\circ\text{C} + 8^\circ\text{C}$$

Dilutions

Answers :

- 1) $[H^+] = 0.50 \text{ M}$ $[Cl^-] = 0.50 \text{ M}$
- 2) $[H^+] = 6.00 \text{ M}$ $[SO_4^{2-}] = 3.00 \text{ M}$
- 3) $[Na^+] = 4.61 \text{ M}$ $[PO_4^{3-}] = 1.54 \text{ M}$
- 4) $[Cu^{2+}] = 0.0801 \text{ M}$ $[SO_4^{2-}] = .0801 \text{ M}$
- 5) a) 0.0625 mol NaOH
b) 0.0625 mol Na^+ 0.0625 mol OH^-
- 6) a) 5.00×10^{-4} mol $CoCl_2$
b) 5.00×10^{-4} mol Co^{2+} 1.00×10^{-3} mol Cl^-
- 7) a) 1.71 mol H_3PO_4
b) 5.12 mol H^+ 1.71 mol PO_4^{3-}
- 8) a) 5.79×10^{-3} mol $Ca(OH)_2$
b) 5.79×10^{-3} mol Ca^{2+} 1.16×10^{-2} mol OH^-
- 9) a) 2.91 M H_2SO_4
b) $[H^+] = 5.81 \text{ M}$ $[SO_4^{2-}] = 2.91 \text{ M}$
- 10) a) 2.44 M HCl
b) $[H^+] = 2.44 \text{ M}$ $[Cl^-] = 2.44 \text{ M}$
- 11) a) 0.667 M KOH
b) $[K^+] = 0.667 \text{ M}$ $[OH^-] = 0.667 \text{ M}$
- 12) a) 0.395 M $LiNO_3$ 1.51 M $FeCl_3$
b) $[Li^+] = 0.395 \text{ M}$ $[NO_3^-] = 0.395 \text{ M}$
 $[Fe^{3+}] = 1.51 \text{ M}$ $[Cl^-] = 4.54 \text{ M}$
- 13) 62.5 mL of 12.0 M HCl mixed with water to raise volume to 250.0 mL
- 14) 2.77 mL of 18.0 M H_2SO_4 mixed with water to raise volume to 50.0 mL
- 15) 3.33 mL of 15.0 M CH_3COOH mixed with water to raise volume to 100.0 mL
- 16) 50.0 mL of 15.0 M NH_3 mixed with water to raise volume to 500.0 mL

Unit 7—Stoichiometry

1. Given the following balanced equation, answer the questions following it:



- a) If 5.5 moles of H_2 are reacted, how many moles of NF_3 will be consumed?

$$5.5 \text{ mols } \text{H}_2 \times \frac{2}{3} =$$

Answer 3.67 mol

- b) In order to produce 0.47 moles of HF , how many moles of NF_3 would be consumed?

$$0.47 \text{ mol} \times \frac{2}{6} =$$

Answer 0.157 mol

- c) If you needed to produce 180.6 g of N_2 , how many moles of H_2 would you need to start with?

$$180.6 \text{ g} \times \frac{1 \text{ mol}}{28 \text{ g}} \times \frac{3}{1} =$$

Answer 19.35 mol

- d) If you completely react 17.04 g of NF_3 , what mass of HF will be produced?

$$17.04 \text{ g} \times \frac{1 \text{ mol}}{71 \text{ g}} \times \frac{6}{2} \times \frac{20.0 \text{ g}}{1 \text{ mol}} =$$

Answer 14.4 g

2. Given the following balanced equation, answer the questions following it:



- a) If 3.56 moles of HBr are reacted, how many Litres of Br_2 will be formed at STP?

$$3.56 \text{ mol} \times \frac{3}{5} \times \frac{22.4 \text{ L}}{1 \text{ mol}} =$$


Answer 47.85 L

- b) In order to produce 3.311×10^{24} molecules of Br_2 , what mass of HBr is needed?

$$3.311 \times 10^{24} \text{ molec} \times \frac{1 \text{ mol}}{6.022 \times 10^{23}} \times \frac{5}{3} \times \frac{80.9 \text{ g}}{1 \text{ mol}} =$$

Answer 741.6 g

4. Given the following balanced equation, answer the questions below it.



- a) In a titration, 18.20 mL of 0.300 M Ba(OH)_2 is required to react completely with a 25.0 mL sample of a solution of HNO_3 . Find the $[\text{HNO}_3]$.

$$\text{mols Ba(OH)}_2 = \frac{0.300 \text{ mols}}{\cancel{L}} \times 0.0182 \cancel{L} = 0.00546$$

$$\text{mols HNO}_3 = 0.00546 \text{ mol} \times \frac{2}{1} = 0.01092$$

$$[\text{HNO}_3] = \frac{0.01092 \text{ mol}}{0.0250 \cancel{L}} = \text{Answer } \underline{0.437 \text{ M}}$$

- b) In a titration, 11.06 mL of 0.200 M HNO_3 is required to react completely with a sample of 0.250 M Ba(OH)_2 . Find the volume of the Ba(OH)_2 sample.


$$\text{mols HNO}_3 = \frac{0.200 \text{ mols}}{\cancel{L}} \times 0.01106 \cancel{L} = 0.002212 \text{ mol}$$

$$\text{mols Ba(OH)}_2 = 0.002212 \times \frac{1}{2} = 0.001106 \text{ mols}$$



$$V = \frac{0.001106 \text{ mols}}{0.250 \text{ M}} = \text{Answer } \underline{0.004424 \text{ L}}$$

5. Given the following balanced equation, answer the questions below it.



- a) If 317.5 grams of Cu are placed into 756.0 grams of HNO_3 , determine which reactant is in excess.

$$\text{Cu } 317.5 \text{ g} \times \frac{1 \text{ mol}}{63.5 \text{ g}} = 5.0 \text{ mol Cu} \times \frac{2}{3} \times \frac{30 \text{ g}}{1 \text{ mol}} = 99.99$$

$$\text{HNO}_3 = 756.0 \text{ g} \times \frac{1 \text{ mol}}{63 \text{ g}} = 12.0 \text{ mol} \times \frac{2}{8} \times \frac{30 \text{ g}}{1 \text{ mol}} = 90$$

Cu in excess

- b) If the reaction in (a) is carried out, what mass of NO will be formed?

~~2 mol HNO₃~~

HNO_3 is limiting reagent:

Answer 90.0 g

Unit 8— Atoms, Periodic Table and Bonding

1. Give the number of protons, neutrons and electrons in the following:

Isotope	Protons	Neutrons	Electrons
$^{194}_{77}\text{Ir}^{3+}$	77	117	74
$^{202}_{80}\text{Hg}^{2+}$	80	122	78
$^{125}_{52}\text{Te}^{2-}$	52	73	54
$^{263}_{106}\text{Sg}$	106	157	106
$^2_1\text{H}^+$	1	1	0

2. Give the nuclear notation of the following:

Isotope	Protons	Neutrons	Electrons
$^{262}_{105}\text{Db}^{+2}$	105	157	103
$^{123}_{51}\text{Sb}^{+3}$	51	72	48
$^{75}_{33}\text{As}^{-3}$	33	42	36
$^{133}_{54}\text{Xe}$	54	79	54
$^{244}_{94}\text{Pu}^{+3}$	94	150	91

3. Write the ground state electron configurations (eg. $1s^2 2s^2 2p^6$) for the following atoms or ions. You may use the core notation.

- a) P $[Ne] 3s^2 3p^3$
b) Mo $[Kr] 5s^2 4d^4$
c) Se $[Ar] 4s^2 3d^{10} 4p^4$
d) Rb $[Kr] 5s^1$
e) Cl^- $[Ne] 3s^2 3p^6$
f) Al^{3+} $[He] 2s^2 2p^6$
g) K^+ $[Ne] 3s^2 3p^6$
h) S^{2-} $[Ne] 3s^2 3p^6$

4. In order to become stable,

an atom of Sr will lose 2 electrons and become the ion Sr^{+2}

an atom of As will gain 3 electrons and become the ion As^{-3}

an atom of Al will lose 3 electrons and become the ion Al^{+3}

an atom of Se will gain 2 electrons and become the ion Se^{-2}

an atom of N will gain 3 electrons and become the ion N^{-3}

an atom of I will gain 1 electrons and become the ion I^-

an atom of Cs will lose 1 electrons and become the ion Cs^+

an atom of Te will gain 2 electrons and become the ion Te^{-2}

Circle the metalloid: Be Rb Os Ge Pb Al

Circle the most reactive element in the following: Na Mg Si Al Ar

Circle the most reactive element in the following: Na K Rb Cs Li

Circle the most reactive element in the following: Cl Br I At Ne

Circle the element with the largest atomic radius of these: Na Mg Si Al Ar

Circle the element with the largest atomic radius of these: N P As Sb Bi

11 Circle the element with the largest ionization energy of these: K Ca Ga As **Kr**

12 Circle the element with the largest ionization energy of these: **C** Si Ge Sn Pb

13 What is meant by ionization energy? **E to remove outer e⁻**

14 Circle the element with the largest density of these: C Si Ge Sn **Pb**

15 Circle the element with the largest density of these: Na K Rb **Cs** Li

16 Circle the element with the highest electronegativity of these: **Mg** Sr Ba Ra

17 Circle the element with the highest electronegativity of these: Mg Si S **Cl**

18 Circle the element with the highest electronegativity of these: **F** Cl Br I


19 What is meant by electronegativity? **attraction of an atom for e⁻ of another atom**

20 Circle the most metallic element of these: Be Mg Ca Sr **Ba**


21 Circle the most metallic element of these: B Al Ga In **Tl**

22 Circle the most metallic element of these: **Ga** Ge Se Br Kr

23 Write a balanced equation for the reaction of potassium with water.



24 Write a balanced equation for the reaction of aluminum with bromine.



25 In an ionic bond, electrons are

- a. shared equally by two atoms
- b. shared unequally by two atoms
- c.** transferred from a metal to a non-metal
- d. transferred from a non-metal to a metal
- e. closer to one end of a molecule, forming a temporary dipole

Answer _____

26 In a covalent bond, electrons are

- f.** shared equally by two atoms
- g. shared unequally by two atoms
- h. transferred from a metal to a non-metal
- i. transferred from a non-metal to a metal
- j. closer to one end of a molecule, forming a temporary dipole

Answer _____

27 In London forces, electrons are

- p. shared equally by two atoms
- q. shared unequally by two atoms
- r. transferred from a metal to a non-metal
- s. transferred from a non-metal to a metal
- t.** closer to one end of a molecule, forming a temporary dipole

Answer _____

28. What physical evidence do we have that ionic bonds are very strong?

Ionic compounds have high melting points




29. Write electron-dot diagrams for:


MgCl₂ (ionic)

PBr₃ (covalent)

SeF₂ (covalent)

CH₃CH₂I (covalent)







Yet More Lewis Structures – Answers

For those of you that enjoy such things, some more Lewis structures to draw:


1) BSF




2) HBr



3) C₂H₅OH (ethanol)



4) N₂F₄



5) SF₆

